



Anti-DYKDDDDK conjugated to SureLight® APC

Product Number D3-1718-1MG
Amount 1 mg total protein
Store at 2-8°C

Form/Shipping & Storage

Supplied lyophilized. Upon receipt, store at -20 °C. Reconstitute with 1 ml of ddH₂O and store at 2-8 °C. Phycobiliproteins are sensitive to freeze-thaw cycles.

Handling

We recommend that the investigator determine the appropriate working concentration for their specific application. Avoid exposure to heat and light.

Buffer


Upon reconstitution, the product is in 10 mM tris (pH 8.2) + 150 mM NaCl + 0.1% BSA + 50 mM sucrose

Stability

Product should be stored at 2-8°C in the dark and be used within 1 year. If further dilution of the conjugate is required, use diluted material within one month.

Note

For research use only, not for diagnostic or therapeutic use.



Fluorescence excitation and emission spectra of APC in 100 mM sodium phosphate (pH 7.2) + 1 mM EDTA and 1 mM sodium azide. The emission scan was taken with excitation at 630 nm. The excitation scan was taken with an emission at 660 nm. The curves were normalized to equalize peak heights.

Spectral Characteristics

Visible absorption maxima	652
Emission maximum	657

References

Puertollano R, Alonso MA. MAL, an integral element of the apical sorting machinery, is an itinerant protein that cycles between the trans-Golgi network and the plasma membrane. *Mol Biol Cell.* 1999 Oct;10(10):3435-3447.

Roy SF, Laporte SA, Escher E, Leduc R, Guillemette G. Epitope tagging and immunoreactivity of the human angiotensin II type 1 receptor. *Can J Physiol Pharmacol.* 1997 Jun;75(6):690-695.



COLUMBIA BIOSCIENCES